

Search for VHE γ -ray emission from the globular cluster M13 with the MAGIC telescope

H. Anderhub^a, L. A. Antonelli^b, P. Antoranz^c, M. Backes^d, C. Baixeras^e, S. Balestra^c, J. A. Barrio^c, D. Bastieri^f, J. Becerra González^g, J. K. Becker^d, W. Bednarek^h, K. Berger^h, E. Bernardiniⁱ, A. Biland^a, R. K. Bock^{j,f}, G. Bonnoli^k, P. Bordas^l, D. Borla Tridon^j, V. Bosch-Ramon^l, D. Bose^c, I. Braun^a, T. Bretz^m, I. Britvitch^a, M. Camara^c, E. Carmona^j, S. Commichau^a, J. L. Contreras^c, J. Cortinaⁿ, M. T. Costado^{g,o}, S. Covino^b, V. Curtef^d, F. Dazzi^{p,***}, A. De Angelis^p, E. De Cea del Pozo^q, R. de los Reyes^c, B. De Lotto^p, M. De Maria^p, F. De Sabata^p, C. Delgado Mendez^{g,*}, A. Dominguez^r, D. Dorner^a, M. Doro^f, D. Elsaesser^m, M. Errandoⁿ, D. Ferenc^s, E. Fernándezⁿ, R. Firpoⁿ, M. V. Fonseca^c, L. Font^e, N. Galante^j, R. J. García López^{g,o}, M. Garczarczykⁿ, M. Gaug^g, F. Goebel^{j,****}, D. Hadasch^e, M. Hayashida^j, A. Herrero^{g,o}, D. Hildebrand^a, D. Höhne-Mönch^m, J. Hose^j, C. C. Hsu^j, T. Jogler^j, D. Kranich^a, A. La Barbera^b, A. Laille^s, E. Leonardo^k, E. Lindfors^t, S. Lombardi^f, F. Longo^p, M. López^f, E. Lorenz^{a,j}, P. Majumdarⁱ, G. Maneva^u, N. Mankuzhiyil^p, K. Mannheim^m, L. Maraschi^b, M. Mariotti^f, M. Martínezⁿ, D. Mazinⁿ, M. Meucci^k, M. Meyer^m, J. M. Miranda^c, R. Mirzoyan^j, H. Miyamoto^j, J. Moldón^l, M. Moles^r, A. Moralejoⁿ, D. Nieto^c, K. Nilsson^t, J. Ninkovic^j, N. Otte^{j,**}, I. Oya^c, R. Paoletti^k, J. M. Paredes^l, M. Pasanen^t, D. Pascoli^f, F. Pauss^a, R. G. Pegna^k, M. A. Perez-Torres^r, M. Persic^{p,v}, L. Peruzzo^f, F. Prada^r, E. Prandini^f, N. Puchadesⁿ, I. Reichardtⁿ, W. Rhode^d, M. Ribó^l, J. Rico^{w,n}, M. Rissi^a, A. Robert^e, S. Rügamer^m, A. Saggion^f, T. Y. Saito^j, M. Salvati^b, M. Sanchez-Conde^r, K. Sataleckaⁱ, V. Scalzotto^f, V. Scapin^p, T. Schweizer^j, M. Shayduk^j, S. N. Shore^x, N. Sidroⁿ, A. Sierpowska-Bartosik^q, A. Sillanpää^t, J. Sitarek^{j,h}, D. Sobczynska^h, F. Spanier^m, A. Stamerra^k, L. S. Stark^a, L. Takalo^t, F. Tavecchio^b, P. Temnikov^u, D. Tesaroⁿ, M. Teshima^j, M. Tluczykontⁱ, D. F. Torres^{w,q}, N. Turini^k, H. Vankov^u, R. M. Wagner^j, W. Wittek^j, V. Zabalza^l, F. Zandanel^r, R. Zaninⁿ, J. Zapatero^e,

ABSTRACT

Based on MAGIC observations from June and July 2007, we have obtained an integral upper limit to the VHE energy emission of the globular cluster M13 of $F(E > 200 \text{ GeV}) < 5.1 \times 10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$, and differential upper limits for $E > 140 \text{ GeV}$. Those limits allow us to constrain the population of millisecond pulsars within M13 and to test models for acceleration of leptons inside their magnetospheres and surrounding. We conclude that in M13 either millisecond pulsars are fewer than expected or they accelerate leptons less efficiently than predicted.

Subject headings: gamma rays: observations, globular cluster: individual (M13)

^aETH Zurich, CH-8093 Switzerland

^bINAF National Institute for Astrophysics, I-00136 Rome, Italy

^cUniversidad Complutense, E-28040 Madrid, Spain

^dTechnische Universität Dortmund, D-44221 Dortmund, Ger-

many

^eUniversitat Autònoma de Barcelona, E-08193 Bellaterra, Spain

^fUniversità di Padova and INFN, I-35131 Padova, Italy

^gInst. de Astrofísica de Canarias, E-38200 La Laguna, Tenerife,

1. Introduction

Globular clusters (GC) are very interesting sites for probing high energy processes due to their large content of evolved objects. Millisecond pulsars (MSP) constitute a large fraction of these objects and it has been estimated that a typical massive GC contains of the order of 100 of them (Tavani 1993). Moreover, the largest sample of MSPs discovered up to now in radio observations are located in the GC Ter 5 (23 MSP), and Tuc 47 (22 MSP) (see e.g. Camilo & Rasio 2005).

Fluxes of TeV γ -rays from GC detectable by current Cherenkov telescopes have been predicted based on estimates on the population of MSPs and the efficiency of lepton acceleration in their surrounding (see Bednarek & Sitarek (2007) and Venter et al. (2009)). These γ -rays would be produced by accelerated leptons scattering off photons of the microwave background radiation or the thermal emission of an extremely dense cluster of solar mass stars inside the GC. Acceleration of leptons could take place in i) the

shocks within the GC, coming from the collision of the winds of MSPs, or ii) in the pulsar inner magnetosphere or their wind regions. In addition, γ -rays in the sub-TeV energy range could also be originated in the inner MSP magnetosphere directly, as it is predicted in the calculations by Bulik et al. (2000); Harding, Usov & Muslinov (2005), or could be produced in the vicinity of radio emitting blocked pulsars (Aharonian et al. 2005; Albert et al. 2006) inside low mass binary systems (Tavani 1991).

GCs have been observed occasionally by Cherenkov Telescopes to probe for this possible VHE γ -ray emission. The few experimental results reported in the literature are upper limits on the emission of M13 by the *WHIPPLE* Collaboration (Hall et al. 2003), M15 by the *VERITAS* Collaboration (LeBohec et al. 2003), and ω Centauri by the *CANGAROO* Collaboration (Kabuki et al. 2007). Very recently, the *Fermi* LAT telescope has detected high energy γ -ray emission ($E > 100$ MeV) from one of the closest and most massive GC, Tuc 47 (Guillemot et al. 2009), and *HESS* has obtained an upper limit of 6.7×10^{-13} ph cm $^{-2}$ s $^{-1}$ for energies $E > 800$ GeV (Aharonian et al. 2009), but given the possible complexity of the emission in the GeV range, it is not possible to establish any connection between these results. This *HESS* result constrains the magnetic field in the pulsar nebula as a function of the number of MSP in the GC for the model by Venter et al. (2009), and in the efficiency of the rotational energy conversion of MSPs into relativistic leptons for the model by Bednarek & Sitarek (2007).

In this paper we report the results of observations with the *MAGIC* telescope of the globular cluster M13, and we discuss the constraints that our results impose to the population of millisecond pulsars and their lepton acceleration efficiency. M13 belongs to the class of normal globular clusters, and has an estimated mass of $6 \times 10^5 M_{\odot}$. It is located in the Northern sky at a distance of ~ 7 kpc. Its core radius is about ~ 1.6 pc, with a half mass radius of ~ 3.05 pc (Harris 1999). By now 5 millisecond pulsars have been detected in M13, with periods ranging between 2 and 10 ms. The aforementioned observation of this cluster in search for VHE emission by the *WHIPPLE* Collaboration (Hall et al. 2003) led to a flux upper limit of 1.08×10^{-11} ph. cm $^{-2}$ s $^{-1}$ at energies $E > 500$ GeV.

Spain

^hUniversity of Łódź, PL-90236 Lodz, Poland

ⁱDeutsches Elektronen-Synchrotron (DESY), D-15738 Zeuthen, Germany

^jMax-Planck-Institut für Physik, D-80805 München, Germany

^kUniversità di Siena, and INFN Pisa, I-53100 Siena, Italy

^lUniversitat de Barcelona (ICC/IEEC), E-08028 Barcelona, Spain

^mUniversität Würzburg, D-97074 Würzburg, Germany

ⁿIFAE, Edifici Cn., Campus UAB, E-08193 Bellaterra, Spain

^oDepto. de Astrofísica, Universidad, E-38206 La Laguna, Tenerife, Spain

^pUniversità di Udine, and INFN Trieste, I-33100 Udine, Italy

^qInstitut de Ciències de l'Espai (IEEC-CSIC), E-08193 Bellaterra, Spain

^rInst. de Astrofísica de Andalucía (CSIC), E-18080 Granada, Spain

^sUniversity of California, Davis, CA-95616-8677, USA

^tTuorla Observatory, University of Turku, FI-21500 Piikkiö, Finland

^uInst. for Nucl. Research and Nucl. Energy, BG-1784 Sofia, Bulgaria

^vINAF/Osservatorio Astronomico and INFN, I-34143 Trieste, Italy

^wICREA, E-08010 Barcelona, Spain

^xUniversità di Pisa, and INFN Pisa, I-56126 Pisa, Italy

^{*}now at: Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), E-28040 Madrid, Spain

^{**}now at: University of California, Santa Cruz, CA 95064, USA

^{***}supported by INFN Padova

^{****}deceased

2. Observations and data analysis

The MAGIC telescope is an Imaging Atmospheric Cherenkov Telescope (IACT) located at the Observatory Roque de los Muchachos on the Canarian Island La Palma. It has an exceptional light detection efficiency provided by the combination of a 17 m diameter mirror and a pixelized camera composed of 576 high quantum efficiency, hemispherical photomultiplier tubes (PMT). This allows MAGIC to reach a standard trigger threshold of ~ 60 GeV. For energies above 150 GeV, angular and energy resolutions of the telescope are ~ 0.1 deg. and $\sim 25\%$ respectively (see Albert et al. 2008a for further details). Besides this, in February 2007 its data acquisition system was upgraded with multiplexed 2 GHz Flash Analog-Digital converters which improved the timing resolution of the recorded shower images. Accordingly the sensitivity of MAGIC improved significantly (Aliu et al. 2009) to 1.6% of the Crab Nebula flux above 270 GeV for 50 hours of observation.

We observed M13 at zenith angles ranging from 8° to 31° between June 12th and July 18th of 2007 in false-source tracking (wobble) mode (Fomin et al. 1994), with two directions at $24'$ distance and opposite sides of the source direction. This technique allows for a reliable estimation of the background with no need of extra observation time. The collected data amount to 20.7 hours after rejecting events affected by unstable hardware or environmental conditions. Besides this, events with a collected charge below 300 photo-electrons were rejected in order to maximize the analysis sensitivity. This selection resulted in a sample with a peak energy of 190 GeV.

Data analysis was carried out using the standard MAGIC analysis and reconstruction software chain, which proceeds in several steps. Initially, a standard calibration of the PMT signal pulses is performed (Albert et al. 2008b). Then, pixels containing no useful information for the shower image reconstruction are discarded by an image cleaning procedure (Aliu et al. 2009). Afterwards event image parameters are calculated (Hillas 1985) using the surviving pixels. In addition to the classical Hillas parameters, two timing parameters are computed, namely: the gradient of the arrival times of the Cherenkov photons along the shower axis; and their arrival time spread over the whole shower. The signal-to-noise maximization is achieved using a multidimensional classification procedure based on the Random Forest (RF)

method (Albert et al. 2008c), where a hadron likeness measure, the so called hadronness, is computed for each event based on the image and time parameters. Moreover, a regression RF trained with a Monte Carlo simulated γ -ray sample is used to estimate the energy on an event by event basis. Finally the angle between the major axis of the shower image ellipse and the source position in the camera, the so called Alpha angle, is used to select γ -ray candidates in the direction of the source. To estimate the residual background, the angle Alpha is also computed with respect to a position symmetric to the source position with respect to the camera center. In what follows, this position used to estimate the residual background is referred as the background region.

Main contributions to the systematic error of our analysis are the uncertainties in the atmospheric transmission, the reflectivity of the mirrors (including losses due to surface roughness) and the light catchers, the photon to photo-electron conversion calibration and the photo-electron collection efficiency in the photomultiplier front-end. A detailed discussion of their contribution to the flux uncertainties can be found in Albert et al. (2008a), where they are estimated to add up to 30% of the measured flux value.

3. Results

Figure 1 shows the obtained distribution of the Alpha angle for the source region and the estimated background. It has been obtained for events surviving a hadronness cut tuned to yield an energy independent γ -ray selection efficiency of 80%, estimated by means of a Monte Carlo simulation. We define the signal region as the smaller interval in Alpha angle that contains the 80% of the γ -rays for each energy bin, estimated using a Monte Carlo simulation. Their lower bounds are at Alpha= 0 and the upper ones are shown in second column of Table 1 for each energy bin. We find a total of -23 ± 57 excess events after background subtraction in this signal region for an energy range extending from 140 GeV to 1.1 TeV. In addition a search for signals in a region of 1 deg of radius around M13 yields no positive detection. We have obtained upper limits to the VHE flux from M13 for different energy bins, shown in Table 1. These have been computed using the Rolke method by Rolke et al. (2005) at a 95% confidence level, and they take into account a 30% of systematic uncertainties in the flux level. The upper limit to the integral flux for energies above

$E = 200$ GeV, assuming a spectral index of 2.6, is $5.1 \times 10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$.

Table 1: Differential upper limits

Energy bin GeV	Upper Alpha cut (deg)	Events	Background events	Excess UL (95% CL)	Flux UL ($\text{cm}^{-2} \text{ s}^{-1} \text{ TeV}^{-1}$)
140 – 200	8	487	517 ± 23	37	7.2×10^{-11}
200 – 280	10	683	681 ± 27	95	5.1×10^{-11}
280 – 400	8	254	242 ± 16	75	2.2×10^{-11}
400 – 560	6	62	73 ± 9	14	2.4×10^{-12}
560 – 790	4	32	27 ± 5	27	2.7×10^{-12}
790 – 1120	4	4	5.7 ± 2.4	5.8	3.7×10^{-13}

4. Comparison with models

In Figure 2 we compare our flux upper limits with the theoretical γ -ray spectra calculated by Bednarek & Sitarek (2007). In this model, leptons are injected into the GC volume according to a power-law spectrum, upon acceleration in the shocks produced in the collisions of the pulsar winds of several MSPs. γ -rays are then produced via Inverse Compton scattering of photons from the microwave background radiation and the thermal radiation arising from the whole GC. Thus, the comparison of our experimental upper limits with the different theoretical gamma-ray spectra allows us to constrain the total power of injected leptons (L_e). For this, we require the theoretical gamma-ray spectra to be lower than all the obtained experimental limits, we take into account the light field of the GC (Bednarek & Sitarek 2007) and assume a distance of 7 pc. The upper limits to L_e are reported in Table 2 for different assumptions of the spectrum of the injected leptons, i.e. for different values of the spectral index α between the minimum energy $E_{\min}$ and the maximum energy defined by the escape of leptons from the shock. In the case of mono-energetic injection of leptons we consider two different energies, 1 TeV and 10 TeV. Assuming characteristic values for the parameters of the MSPs in globular clusters (surface magnetic field 10^9 G and rotational period 4 ms), we can translate the limits to L_e into limits to the product of the required number of the MSPs in M13 (N_{MSP}) times the efficiency of the rotational energy conversion of MSPs into relativistic leptons (η), shown in Table 2. For example, in the case of M13, Tavani (1993) predicts the existence of 100 MSPs. On the other hand, the efficiency of lepton injection from the inner magnetospheres of millisecond pulsars has been estimated to be $\eta \sim 0.1$ in terms of the extended polar gap model by Muslinov & Harding (1997). Therefore,




Fig. 1.— Distribution of Alpha for the selected γ -ray candidates from the source (black dots) and the background (histogram) regions.




Fig. 2.— The MAGIC γ -ray flux upper limits for M13 compared with spectra expected for the range of parameters of the model shown in Figure 9 and 10 of Bednarek & Sitarek (2007). The specific γ -ray spectra are calculated for lepton upper energy cut-off at 3 TeV and lower energy cut-off at 1 GeV (black thick) and 100 GeV (black thin), and power-law spectral indices of 2.1 (solid) and 3 (dashed). The γ -ray spectra produced by mono-energetic leptons of 10 TeV and 1 TeV are shown by a grey solid curve and a dashed one respectively. All calculations are computed assuming the conservative value of 1 for the free parameter of the model $N_{\text{MSP}} \cdot \eta$. The Whipple differential upper limit shown here has been derived from the integral quoted in Hall et al. (2003) assuming a spectral index of 2.6.

the likely value of the product, $N_{\text{MSP}} \cdot \eta$, should be of the order of ~ 10 . We show in the corresponding row of Table 2 our estimate of the upper limit to this product for different models of injected spectra of leptons. Moreover, our limits in this product are at the same level than the recently published ones by the *HESS* collaboration making use of data from 47 Tuc (Aharonian et al. 2009). Figure 3 shows these limits in the N_{MSP}, η plane, such that for each set of model parameters the area above the corresponding curve is excluded at 95% CL. For most of the considered models $N_{\text{MSP}} \cdot \eta$ is significantly below ~ 10 . The only exception is the model with the steep spectrum of leptons which extends down to 1 GeV. Note that even if the number of MSP in M13 is only equal to 5 (as presently observed, Camilo & Rasio 2005), we can already obtain the acceleration efficiency of leptons to be ~ 0.1 in the case of their injection with the hard (spectral index 2.1) and mono-energetic spectrum respectively (see Figure 3).

The γ -ray spectra produced in the curvature process inside the inner pulsar magnetospheres are predicted to cut-off below ~ 100 GeV (Bulik et al. 2000; Harding, Usov & Muslinov 2005). Thus, they can not extend to the energy region investigated by our measurement. An Inverse Compton γ -ray component is expected from leptons accelerated in the inner magnetospheres which extends > 100 GeV. But its flux is predicted to be at the level of several orders of magnitude below the here presented upper limits for most of the energy range covered by our limits. For example, Venter et al. (2009) computation of VHE spectrum for GC Tuc 47 and Ter 5 predicts a flux level similar to the ones of the model by Bednarek & Sitarek (2007) for these pulsars only for narrow energy band above $E = 1$ TeV.

Therefore, we conclude that the inner magnetosphere γ -ray emission of millisecond pulsars is not likely to be detected by present observations with an analysis threshold of the order of ~ 100 GeV even

from globular clusters containing hundreds of MSPs.

5. Conclusions

We have obtained the strongest upper limits to date on the VHE γ -ray flux from the massive globular cluster M13. Our upper limit is ~ 2 times lower than the previous limit for VHE energy emission from M13 quoted by *WHIPPLE*, and extends to energies down to 140 GeV. Our upper limits allow us to constrain the population of the millisecond pulsars expected in M13 and the acceleration scenarios of leptons by millisecond pulsars. Our result strongly suggests that either the number of millisecond pulsars in M13 is significantly lower than the estimate of ~ 100 , or the energy conversion efficiency from millisecond pulsars to relativistic leptons is significantly below the value quoted in recent modeling of high energy processes in the magnetospheres of millisecond pulsars. Our upper limits regarding Bednarek & Sitarek (2007) model parameters are the same level than the ones obtained by the *HESS* collaboration making use of observations of Tuc 47.

Acknowledgements

We would like to thank the Instituto de Astrofísica de Canarias for the excellent working conditions at the Observatorio del Roque de los Muchachos in La Palma. The support of the German BMBF and MPG, the Italian INFN and Spanish MICINN is gratefully acknowledged. This work was also supported by ETH Research Grant TH 34/043, by the Polish MNiSzW Grant N N203 390834, and by the YIP of the Helmholtz Gemeinschaft.

REFERENCES

- Aharonian, F. et al. 2005, *A&A*, 442, 1
- Aharonian, F. et al. 2009, arXiv:0904.0361v1 [astro-ph]
- Albert, J. et al. 2006, *Science*, 312, 1771
- Albert, J. et al. 2008a *ApJ* 674, 1037
- Albert, J. et al. 2008b, *Nucl. Instrum. Methods A*, 594, 407
- Albert, J. et al. 2008c, *Nucl. Instrum. Methods A*, 588, 424
- Aliu, E. et al. 2009 *Astropart. Phys.* 30, 293

Table 2: Upper limits on the power of injected leptons (L_e) and in $N_{\text{MSP}} \cdot \eta$

E_{min} α	100 GeV 2.1	100 GeV 3.0	1 GeV 2.1	1 GeV 3.0	mono: 1 TeV	mono: 10 TeV
L_e ($\times 10^{35}$ erg s $^{-1}$)	0.6	1.0	1.0	60	0.2	0.5
$N_{\text{MSP}} \cdot \eta$	0.5	1.0	1.0	50	0.2	0.4

Bednarek, W. & Sitarek, J. 2007 MNRAS 377, 920 (BS07)

Bulik, T., Rudak, B. & Dysk, J. 2000 MNRAS 317, 97

Camilo, F. & Rasio, F.A. 2005 Proc. Binary Radio Pulsars, ASP Conf. Ser. Vol. 328, eds. F.A. Rasio & I.H. Stairs (San Francisco: ASP), p. 147

Fomin, V. P., Stepanian, A., Lamb, R. C., Lewis, D. A., Punch, M. & Weekes, T. C. 1994 Astropart. Phys. 2, 137

Guillemot, L. et al. 2009 Rencontres de Moriond (2009)

Hall, T.A., Bond, I.H., Bradbury, S.M. et al. 2003 ApJ 583, 853

Harding, A.K., Usov, V.V. & Muslimov, A. 2005 ApJ 622, 531

Harris, W.E. 1991 Ap&SS 267, 95

Hillas, A. M. 1985 Proc. of the 19th ICRC (La Jolla)

Kabuki, S., Enomoto, R., Bicknell, G.V. et al. 2007 ApJ 668, 968

LeBohec, S., Baltz, E.A., Bond, I.H. et al. 2003 Proc. 28th ICRC (Tsukuba)

Muslimov, A.G. & Harding, A.K. 1997 ApJ 485, 735

Rolke, W., Lopez, A., Conrad, J. & James, F. 2005 Nucl.Instrum.Meth.A 551, 493

Tavani, M. 1991 ApJ 379, L69

Tavani, M. 1993 ApJ 407, 135

Venter, C., de Jager, O.C., & Clapson, A.C. 2009, arXiv:0903.3031 [astro-ph]




Fig. 3.— Exclusion contours in the N_{MSP}, η plane for the different models considered in the text. The model with parameters $\alpha=3$ and cut-off at 100 GeV overlaps the $\alpha=2.1$ with cut-off at 1 GeV one. The horizontal and vertical long-dashed black lines show the reference values of $\eta=0.1$ and $N_{MSP}=5$ respectively.